

| PATIENT INFORMATION  |  |
|--|--|
| Patient Name: _____  | Date: _____  |
| Address: _____   | Phone: _____   |
| City, State, Zip: _____  |  |
| DOB: _____ Weight: _____ Height: _____   | <b>Diabetic (Circle one):</b> Yes / No   |
| Diagnosis / ICD 10 Code: _____   | Allergies: _____   |
| <input type="checkbox"/> INITIATION OF THERAPY   | <input type="checkbox"/> CHANGE OF INTERVAL  |
| <input type="checkbox"/> CHANGE OF PROVIDER  |  |
| If the patient cannot provide recent PPD results (within 12 months), then perform PPD.   |  |
| <b>TUBERCULOSIS SCREENING:</b> <input type="checkbox"/> PPD <b>Date Performed:</b> _____ <b>Date Read:</b> _____   |  |
| <input type="checkbox"/> PPD results provided by patient per Dr. _____ <b>Date:</b> _____  |  |
| <b>Result:</b> <input type="checkbox"/> <b>Negative</b> → May Initiate Remicade® Therapy   |  |
| <input type="checkbox"/> <b>Positive</b> → <input type="checkbox"/> Chest X-Ray performed <b>Date Performed:</b> _____ <input type="checkbox"/> <b>Negative result</b> → May initiate Remicade®<br><input type="checkbox"/> <b>Positive result</b> → TB treatment initiated  |  |
| <input type="checkbox"/> RN to teach disease management  |  |
| <b>Remicade® - Choose Regimen:</b> <input type="checkbox"/> <b>Remicade</b> <input type="checkbox"/> <b>Avsola</b> <input type="checkbox"/> <b>Inflectra</b> <input type="checkbox"/> <b>Renflexis</b>   |  |
| <input type="checkbox"/> 3 mg/kg _____ mg/NS 0.9% 250 mL or 500 mL of normal saline IV over 2 hours  | <input type="checkbox"/> NS 0.9% 50 mL IVPB to flush Remicade from IV tubing                             |
| <input type="checkbox"/> 5 mg/kg _____ mg/NS 0.9% 250 mL or 500 mL of normal saline IV over 2 hours  | <input type="checkbox"/> Concomitant po Methotrexate _____ mg/week<br>(Methotrexate RX given to Patient) |
| <input type="checkbox"/> 10 mg/kg _____ mg/NS 0.9% 250 mL IV over 2 hours  | <input type="checkbox"/> Infuse at weeks 0,2, and 6 then every 8 weeks                                   |
| <input type="checkbox"/> _____ mg/kg _____ mg/NS 0.9% 250 mL or 500 mL of normal saline IV over 2 hours  | <input type="checkbox"/> Other frequency: _____  |
| <input type="checkbox"/> Repeat dose in 2 weeks and then every 6 weeks   |  |
| <b>Duration of Therapy:</b> <input type="checkbox"/> x 1 year  |  |
| <input type="checkbox"/> Other: _____  |  |
| * Start infusion at 10 mL/hr. Double the rate after 15 minutes (as tolerated) for the first hour of infusion. (After first 15 minutes → 20 mL/hr, at 30 minutes → 40 mL/hr, at 45 minutes → 80 mL/hr, at 1 hr → 160 mL/hr). At 90 minutes into the infusion, the rate may be increased for the last time to a maximum rate of 250 mL/hr. |  |
| <b>Pre-Medications:</b>  |  |
| <input type="checkbox"/> <b>Diphenhydramine</b> _____ mg, po -or- IV (circle one), prior to start of infusion  |  |
| <input type="checkbox"/> <b>Acetaminophen</b> 650 mg po prior to start of infusion   |  |
| <input type="checkbox"/> <b>Prednisone</b> _____ mg, po -or- <input type="checkbox"/> Methylprednisolone 40 mg IVP -or- <input type="checkbox"/> Hydrocortisone 100 mg IVP   |  |
| <input type="checkbox"/> <b>Other:</b> _____   |  |
| <b>* Administer pre-medications 15 to 30 minutes prior to start of infusion *</b>  |  |
| <b>IV Access:</b>  |  |
| <input type="checkbox"/> Start PIV if no IV access available <input type="checkbox"/> Maintain current central line access   |  |
| <b>Catheter Care:</b>  |  |
| <input type="checkbox"/> Sodium Chloride 0.9% _____ mL IV before and after each IV access and PRN per protocol   |  |
| <input type="checkbox"/> Sodium Chloride 0.9% _____ mL as above <b>AND</b> Heparin 100 Units /mL _____ mL IV flush after second saline flush and PRN   |  |
| * Dressing changes weekly and PRN                      * Antimicrobial dressing PRN  |  |
| * May obtain blood from IV access for labs                      * May use Cathflo 2 mg/2 mL sterile water IVP 2 mL per lumen; May repeat after 2 hours' x 1  |  |
| <b>Labs:</b>   |  |
| <input type="checkbox"/> CBC q _____ <input type="checkbox"/> CMP q _____ <input type="checkbox"/> CRP q _____ <input type="checkbox"/> ESR q _____ <input type="checkbox"/> LFTs q _____ <input type="checkbox"/> X-ray _____ <input type="checkbox"/> Other _____  |  |
| <b>Standard Orders for Side Effects:</b>   |  |
| <input type="checkbox"/> <b>Promethazine</b> 25 mg - 1-2 tabs po q 4-6 hrs PRN nausea / vomiting   | <input type="checkbox"/> <b>Diphenhydramine</b> 25 mg - 1 to 2 caps po PRN                               |
| <input type="checkbox"/> <b>Acetaminophen</b> 325 mg - 2 tabs po q 4-6 hrs PRN HA, myalgia, fever  | <input type="checkbox"/> <b>Diphenhydramine</b> 25 mg -or- 50 mg IV x 1 dose PRN                         |
| <input type="checkbox"/> <b>Promethazine</b> 25 mg IV/IM x 1 dose PRN nausea / vomiting  | <input type="checkbox"/> <b>Other:</b> _____   |
| <b>Monitoring Parameters:</b>  |  |
| <input type="checkbox"/> Obtain vital signs and temperature q 15 mins for the 1 <sup>st</sup> hour, then q 30 mins for the remainder of the infusion   |  |
| <input type="checkbox"/> Observe patient for 30 mins following the infusion  |  |
| <input type="checkbox"/> Instruct patient to report symptoms of chills, fever, headache, sore throat, pain, etc.   |  |
| <input type="checkbox"/> Other: _____  |  |
| <b>Anaphylactic Reaction (AR):</b>   |  |
| <b>Epinephrine</b> (based on patient weight)   |  |
| EpiPen® Auto-injector 0.3 mg (1:1000) - Inject IM -or- SubQ to patients who weigh ≥ 66 lbs (≥ 30 kg); may repeat in 3-5 mins x 1 if necessary  |  |
| EpiPen Jr® Auto-injector 0.15mg (1:2000) - Inject IM -or- SubQ to patients who weigh 33 - 66 lbs (15-30 kg); may repeat in 3-5 mins x 1 if necessary   |  |
| <b>Diphenhydramine</b> 50mg (1mL) - Give 50 mg slow IVP, administer IM if no IV access; may repeat x 1 after 10 mins, if necessary   |  |
| <b>Hydrocortisone</b> 100mg - Give 100 mg IVP -or- IM if no IV access  |  |
| <b>Sodium Chloride</b> 0.9% 500 mL infuse IV at a rate of 30 mL/hr   |  |
| <b>Other Orders:</b> _____   |  |

Prescribing Physician: \_\_\_\_\_ Address: \_\_\_\_\_ Phone: \_\_\_\_\_

**PHYSICIAN'S SIGNATURE:** \_\_\_\_\_ **DATE:** \_\_\_\_\_

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation.

If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited.

If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.