

## Uplizna® - Inebilizumab Injection

Phone: (877) 246-9104 Fax: (800) 783-9146 www.biotekrx.com

PATIENT INFORMAT	ION (Complete or fax	c existing chart)	PRESCRIBER INFORMA			
Patient Name:			Prescriber Name:			
Address:			State License: NPI #:			
City, State, Zip:			DEA: Phone:			
Phone: 2 <sup>nd</sup> Phone:			Address: Fax:			
DOB: Gender: U Male U Female			City, State, Zip:			
Weight: Ht: Date:			Contact Person:			
E-mail address:			Phone:			
INSURANCE INFORI	MATION: Copy and a	attach the front and	I back of insurance and pres	cription card(s)		
Primary Insurance:			RX Card (PBM):			
City, State, Zip:		BIN: PCN:				
Member ID #: Phone:		City, State, Zip:				
Plan #: Group #:		Plan #:	Group #:			
DIAGNOSIS / CLINIC	AL INFORMATION					
Primary ICD-10 code:	Diagr	nosis 🗆 G36.0 Neurom	yelitis optica			
Is the patient anti-aquaporin-4 antibody positive? ☐ Yes ☐ No ☐ Test pending						
Prior NSMOD therapies trie		·	·			
Hep B vaccination:   Yes		Does the p	atient have active Hepatitis B infec	tion? ☐ Yes ☐ No		
Hepatitis B screening: ☐ H			•			
☐ HB core antibody <b>HBcA</b> k	•		inve - Negative Date.			
			Tubana da isana mina Danis	in Dispeting D	-4	
Does the patient have active or latent TB infection?						
		lote: Uplizna loading de	oses must be administered in a cor	ntrolled setting.		
Expected date of first/next i	nfusion:					
$\ \square$ NKDA $\ \square$ Known drug al	lergies:					
Concurrent Meds:						
PRESCRIPTION / ADMI	NISTRATION					
	NISTRATION Dose	Directions			Quantity / Refills	
	Dose		250mL of 0.9% NS.		Quantity / Refills  ☐ 6 vials - No refills	
Medication  ☐ Uplizna®	Dose 100mg/10mL SDV	Infusion 1: 300mg in 2				
Medication  ☐ Uplizna®  (inebilizumab injection)	Dose 100mg/10mL SDV Each dose 300mg/30mL	Infusion 1: 300mg in 2 Infusion 2: (2 weeks l	ater): 300mg in 250mL of 0.9% NS	S.		
Medication  Uplizna® (inebilizumab injection) Initial dose (two infusions)	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9%	Infusion 1: 300mg in 2 Infusion 2: (2 weeks l Start infusion at 42mL	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes,	S. increase to 125mL		
Medication  ☐ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection	Infusion 1: 300mg in 2 Infusion 2: (2 weeks l Start infusion at 42mL per hour for the next 3	ater): 300mg in 250mL of 0.9% NS	S. increase to 125mL		
Medication  ☐ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of	Infusion 1: 300mg in 2 Infusion 2: (2 weeks last infusion at 42mL per hour for the next 3 finished.	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m	S. increase to 125mL		
Medication  ☐ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection	Infusion 1: 300mg in 2 Infusion 2: (2 weeks la Start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or la	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger	S. increase to 125mL L per hour until		
Medication  ☐ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of	Infusion 1: 300mg in 2 Infusion 2: (2 weeks last infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or last Monitor patient for at	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m	S. increase to 125mL L per hour until		
Medication  ☐ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL	Infusion 1: 300mg in 2 Infusion 2: (2 weeks last infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or last Monitor patient for at reaction.	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion complete.	increase to 125mL L per hour until	□ 6 vials - No refills	
Medication  ☐ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV	Infusion 1: 300mg in 2 Infusion 2: (2 weeks last infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or last Monitor patient for at reaction. Every 6 months (from	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion completing infusion infuse 300mg in 250	increase to 125mL L per hour until letion for infusion		
Medication  ☐ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL	Infusion 1: 300mg in 2 Infusion 2: (2 weeks la Start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or la Monitor patient for at reaction. Every 6 months (from Start infusion at 42mL	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion completes infusion infuse 300mg in 250 per hour for the first 30 minutes,	increase to 125mL L per hour until letion for infusion 0mL of 0.9% NS. increase to 125mL	□ 6 vials - No refills	
Medication  Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  Uplizna® (inebilizumab injection)	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9%	Infusion 1: 300mg in 2 Infusion 2: (2 weeks lands and 2 mL per hour for the next 3 finished. Duration: 2 hours or lands and 2 mL Every 6 months (from Start infusion at 42mL per hour for the next 3	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion completing infusion infuse 300mg in 250	increase to 125mL L per hour until letion for infusion 0mL of 0.9% NS. increase to 125mL	☐ 6 vials - No refills ☐ 3 vials	
Medication  Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  Uplizna® (inebilizumab injection) Maintenance dose (one	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection	Infusion 1: 300mg in 2 Infusion 2: (2 weeks la Start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or la Monitor patient for at reaction. Every 6 months (from Start infusion at 42mL	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion completes infusion infuse 300mg in 250 per hour for the first 30 minutes,	increase to 125mL L per hour until letion for infusion 0mL of 0.9% NS. increase to 125mL	☐ 6 vials - No refills ☐ 3 vials	
Medication  Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  Uplizna® (inebilizumab injection)	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of	Infusion 1: 300mg in 2 Infusion 2: (2 weeks lands and 2 mL per hour for the next 3 finished. Duration: 2 hours or lands and 2 mL Every 6 months (from Start infusion at 42mL per hour for the next 3	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion completifirst infusion) infuse 300mg in 250 per hour for the first 30 minutes, 30 minutes, then increase to 333m	increase to 125mL L per hour until letion for infusion 0mL of 0.9% NS. increase to 125mL	☐ 6 vials - No refills ☐ 3 vials	
Medication  Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  Uplizna® (inebilizumab injection) Maintenance dose (one	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection	Infusion 1: 300mg in 2 Infusion 2: (2 weeks less start infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or less Monitor patient for at reaction.  Every 6 months (from Start infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or less bursting in the start infusion at 42mL per hour for the next 3 finished.	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion completifirst infusion) infuse 300mg in 250 per hour for the first 30 minutes, 30 minutes, then increase to 333m	increase to 125mL L per hour until letion for infusion 0mL of 0.9% NS. increase to 125mL L per hour until	☐ 6 vials - No refills ☐ 3 vials	
Medication  Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  Uplizna® (inebilizumab injection) Maintenance dose (one	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of	Infusion 1: 300mg in 2 Infusion 2: (2 weeks less start infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or less Monitor patient for at reaction.  Every 6 months (from Start infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or less bursting in the start infusion at 42mL per hour for the next 3 finished.	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion completing infusion infuse 300mg in 250 per hour for the first 30 minutes, 30 minutes, then increase to 333m onger	increase to 125mL L per hour until letion for infusion 0mL of 0.9% NS. increase to 125mL L per hour until	☐ 6 vials - No refills ☐ 3 vials	
Medication  Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  Uplizna® (inebilizumab injection) Maintenance dose (one	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL	Infusion 1: 300mg in 2 Infusion 2: (2 weeks last infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or last reaction.  Every 6 months (from Start infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or last infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or last infusion at 4 finished.	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi first infusion) infuse 300mg in 250 per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi least one hour after infusion compi	increase to 125mL L per hour until letion for infusion 0mL of 0.9% NS. increase to 125mL L per hour until	☐ 6 vials - No refills ☐ 3 vials	
Medication  Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  Uplizna® (inebilizumab injection)  Maintenance dose (one infusion)	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  dministered via pump and	Infusion 1: 300mg in 2 Infusion 2: (2 weeks land start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or land start infusion at 42mL per hour for the next 3 finished. Every 6 months (from Start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or land start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or land start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or land start infusion at 42mL per hour for the next 3 finished.	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi first infusion) infuse 300mg in 250 per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi least one hour after infusion compi	increase to 125mL L per hour until letion for infusion 0mL of 0.9% NS. increase to 125mL L per hour until	☐ 6 vials - No refills ☐ 3 vials	
Medication  ☐ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  ☐ Uplizna® (inebilizumab injection)  Maintenance dose (one infusion)  All Uplizna® orders to be a	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  dministered via pump and	Infusion 1: 300mg in 2 Infusion 2: (2 weeks land start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or land start infusion at 42mL per hour for the next 3 finished. Every 6 months (from Start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or land start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or land start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or land start infusion at 42mL per hour for the next 3 finished.	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi first infusion) infuse 300mg in 250 per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi least one hour after infusion compi	increase to 125mL L per hour until letion for infusion  OmL of 0.9% NS. increase to 125mL L per hour until letion for infusion	☐ 6 vials - No refills ☐ 3 vials	
Medication  ☐ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  ☐ Uplizna® (inebilizumab injection)  Maintenance dose (one infusion)  All Uplizna® orders to be an Additional Medication  Premedication Orders  Acetaminophen 650mg F	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  dministered via pump and and Supplies for Hom PO 30 min prior to infusi	Infusion 1: 300mg in 2 Infusion 2: (2 weeks land Infusion 2: (2 weeks land Infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land Infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land Infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land Infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land Infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land Infusion at 42mL peripheral line unless on the Infusion	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi first infusion) infuse 300mg in 250 per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi least one hour after infusion compi	increase to 125mL L per hour until letion for infusion  OmL of 0.9% NS. increase to 125mL L per hour until letion for infusion	□ 6 vials - No refills □ 3 vials Refills: □ 0 □ 1	
Medication  ☐ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  ☐ Uplizna® (inebilizumab injection)  Maintenance dose (one infusion)  All Uplizna® orders to be an Additional Medication  Premedication Orders  Acetaminophen 650mg finfusion; Methylpredniso	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  dministered via pump and and Supplies for Hom PO 30 min prior to infusi	Infusion 1: 300mg in 2 Infusion 2: (2 weeks land Infusion 2: (2 weeks land Infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land Infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land Infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land Infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land Infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land Infusion at 42mL peripheral line unless on the Infusion	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi first infusion) infuse 300mg in 250 per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi otherwise instructed.	increase to 125mL L per hour until  letion for infusion  OmL of 0.9% NS. increase to 125mL L per hour until  letion for infusion  Send quantity suffinfusion,	☐ 6 vials - No refills ☐ 3 vials Refills: ☐ 0 ☐ 1	
Medication  □ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  □ Uplizna® (inebilizumab injection)  Maintenance dose (one infusion)  All Uplizna® orders to be an Additional Medication  Premedication Orders Acetaminophen 650mg Finfusion; Methylpredniso □ Other:	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  dministered via pump and and Supplies for Hom PO 30 min prior to infusi lone 100mg IV 30 min p	Infusion 1: 300mg in 2 Infusion 2: (2 weeks land Start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or land Monitor patient for at reaction.  Every 6 months (from Start infusion at 42mL per hour for the next 3 finished. Duration: 2 hours or land Monitor patient for at reaction.  peripheral line unless on the Infusion on; Diphenhydraminarior to infusion.	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi first infusion) infuse 300mg in 250 per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi otherwise instructed.	increase to 125mL L per hour until  letion for infusion  OmL of 0.9% NS. increase to 125mL L per hour until  letion for infusion  Send quantity suffinfusion,  All caregivers and	☐ 6 vials - No refills ☐ 3 vials Refills: ☐ 0 ☐ 1 icient for medication ancillaries to be given	
Medication  □ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  □ Uplizna® (inebilizumab injection)  Maintenance dose (one infusion)  All Uplizna® orders to be an Additional Medication  Premedication Orders  Acetaminophen 650mg finfusion; Methylpredniso □ Other:  Fluids for Reconstitution	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  dministered via pump and and Supplies for Hom PO 30 min prior to infusi lone 100mg IV 30 min p	Infusion 1: 300mg in 2 Infusion 2: (2 weeks land infusion 2: (2 weeks land infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land infusion at 42mL per hour for the next 3 finished.  Duration: 2 hours or land infusion.	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi first infusion) infuse 300mg in 250 per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion compi otherwise instructed.	increase to 125mL L per hour until letion for infusion  OmL of 0.9% NS. increase to 125mL L per hour until letion for infusion  Send quantity suffinfusion, All caregivers and per protocol from	□ 6 vials - No refills □ 3 vials Refills: □ 0 □ 1  icient for medication ancillaries to be given product package insert.	
Medication  □ Uplizna® (inebilizumab injection) Initial dose (two infusions) Note: Loading doses must be administered in a controlled infusion site.  □ Uplizna® (inebilizumab injection)  Maintenance dose (one infusion)  All Uplizna® orders to be an Additional Medication  Premedication Orders Acetaminophen 650mg Finfusion; Methylpredniso □ Other:	Dose 100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  100mg/10mL SDV Each dose 300mg/30mL diluted in 250mL of 0.9% sodium chloride injection for final concentration of 1.1mg/mL  dministered via pump and and Supplies for Hom PO 30 min prior to infusi lone 100mg IV 30 min p on and Administration itial dose); 0.9% NaCl 2	Infusion 1: 300mg in 2 Infusion 2: (2 weeks land start infusion at 42mL per hour for the next 2 finished. Duration: 2 hours or land Monitor patient for at reaction.  Every 6 months (from Start infusion at 42mL per hour for the next 2 finished. Duration: 2 hours or land Monitor patient for at reaction.  peripheral line unless one Infusion  on; Diphenhydraminarior to infusion.	ater): 300mg in 250mL of 0.9% NS per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion completes infusion infuse 300mg in 250 per hour for the first 30 minutes, 30 minutes, then increase to 333m onger least one hour after infusion completes one hour after infusion completes one hour after infusion completes instructed.	increase to 125mL L per hour until  letion for infusion  OmL of 0.9% NS. increase to 125mL L per hour until  letion for infusion  Send quantity suffinfusion,  All caregivers and per protocol from If patient requires	☐ 6 vials - No refills ☐ 3 vials Refills: ☐ 0 ☐ 1 icient for medication ancillaries to be given	

0.9% NACL 50mL						
0.9% NACL 100mL						
Hypersensitivity / Anaphylaxis Orders*						
In the event of anaphylactic reaction, stop infusion of drug immediately. Start NS 15mL/hour; 0.9% NS 100mL.						
Medicate with epinephrine pen auto-injector 0.3mg/0.3mL IM as needed for anaphylaxis. Call <b>911</b> , physician, or paramedic.						
I authorize ancillary supplies or medical equipment necessary such as needles, syringes, etc. to administer the therapy as needed for administration.						
☐ Skilled nursing visit as needed to establish venous access, administer medication, and assess general status and response to therapy.  *If nursing services will be required for therapy administration, the home health nurse will call for additional orders per state regulations.						
SIGNATURE						
X(Product Substitution Permitted)	Date:					

**CONFIDENTIALITY STATEMENT:** This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation.

If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited.

If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.

Fax completed form to: (800) 783-9146 - Thank you for using BioTek reMEDys®

Rev 5/2022 Page **2** of **2**